

Colorado Insect of Interest

Dobsonfly (Texas Dobsonfly)

Scientific Name: *Corydalus texanus* Banks

Order: Neuroptera/Megaloptera (Lacewings, Antlions, Snakeflies, Dobsonflies and relatives)

Family: Corydalidae (Dobsonflies)

Identification and Descriptive Features: Dobsonflies are impressively large (7-11 cm) flying insects with gray, highly veined membranous wings that fold over their back. Adults also possess prominent jaws, which are greatly enlarged in the males. Immature stages (hellgrammites), found in water, similarly have prominent jaws and an elongate body form with fleshy projection along the sides of the abdomen.

Distribution in Colorado: This species is associated with larger rivers. In Colorado, dobsonflies are most commonly noted in Mesa County near the Colorado River. However, they occur in parts of all the major western Colorado river systems (Green, White, San Miguel, Gunnison) and are also found in the Purgatoire River as it runs through Las Animas and Otero counties.

Life History and Habits: Adult dobsonflies emerge beginning in early June and are present through midsummer, with peak numbers usually found in July. The sexes are easily determined because of substantial differences in the mouthparts. Those of the females are shorter but with curved cutting edges and they are capable of producing a painful pinch. Jaws of the males are substantially elongated and thinner, projecting forward from the head. These are apparently used in stylized mating behaviors and during fights for defense of territory, when the jaws are used to scoop or flip rival males. However, despite their fearsome appearance the jaws can not be used for feeding nor are they strongly muscled. Both sexes may be attracted to outdoor lights at night. Males tend to disperse longer distances from the riverways.



Figure 2. Dobsonfly egg mass laid on a leaf overhanging a river.



Figure 1. The dobsonfly *Corydalus texanus*, collected from Mesa County. The male is on the left, female on the right.



Figure 3. Hellgrammite, the larva of a dobsonfly.

Eggs are laid in white masses, about the size of a quarter, usually on vegetation hanging over the streams. The under surfaces of bridges are also particularly good spots to find egg masses. Larvae hatching from the eggs drop into the water and begin their development.

The immature form of a dobsonfly develops in water and is known as a *hellgrammite*. These are top predators, feeding on larvae of caddisflies, black flies and other aquatic insects. (They are also prized as a fish bait.)

After a year of development - or perhaps two years - the larva leaves the river and builds a pupation chamber under rocks or logs near stream margins. Adults hatch after a few weeks and have a short life span.